

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

REMARKS

Amendments are made to the specification.

In this preliminary amendment, SEQ. ID. NO. 12 is changed to SEQ. ID. NO. 3, and SEQ ID. NO. 25 is changed to SEQ ID. NO. 4. In addition, the references to SEQ. ID. NOs. 3-11, 12-24 and 26-79 are deleted from the specification. References to SEQ. ID. NOs. 3-11, 12-24 and 26-79 have been replaced by descriptions found originally on pages 55-59 of the specification.

These amendments represent only a renumbering of these sequences so that the Sequence Listing will have sequence IDs with consecutive numbers. That is, SEQ. ID. NOs. 1 through 4 are listed in the sequence listing, corresponding to original SEQ. ID. NOs. 1, 2, 12 and 25, respectively.

Applicants also note that, in the sequence listing filed concurrently with this Preliminary Amendment, sequences are listed for SEQ. ID. NOs. 1 through 4, that were not explicitly listed in the original specification. However, these are supported because the sequences were known in the art and referenced in the specification. Support for these listed sequences is as follows:

Support for SEQ. ID. NO. 1 is found in the specification on page 7, lines 32 to 35, with reference to United States Patent No. 5,489,430.

Support for SEQ. ID. NO. 2 is found in the specification on page 8, line 1, referring to GeneBank Accession No. AB023292.

Support for SEQ ID. NO. 3 (renumbered from SEQ. ID. NO. 12 in the amendment) is found on page 20, lines 13 to 19, referring to WO97/36924.

Support for SEQ ID. NO. 4 (renumbered from SEQ. ID. NO. 25 in the amendment) is found on page

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

22, lines 8 to 10, referring to GeneBank Accession No. AB023292.


Applicants therefore assert that no new matter is added by this Preliminary Amendment.

Attached hereto is a marked-up version of the changes made to the by the current amendment. The attached page is captioned "Version with markings to show changes made."

In the event that any fees are due in connection with this paper, please charge our Deposit Account No. 01-2340.

Respectfully submitted,

ARMSTRONG, WESTERMAN & HATTORI, LLP


Daniel A. Geselowitz, Ph.D.
Agent for Applicants
Reg. No. 42,573

DAG/plb
Atty. Docket No. **010898**
Suite 1000, 1725 K Street,
N.W.
Washington, D.C. 20006

(202) 659-2930



23850

PATENT TRADEMARK OFFICE

Enclosures: Version with markings to show changes made

H:\FLOATERS\DAG\Amendments\010898 pre. amend 3-11-03

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE SPECIFICATION:

Please amend the specification as follows:

Please replace the paragraph beginning on page 12, line 10, with the following rewritten paragraph:

As the gene region nonessential for the growth of herpesvirus in the case of Marek's disease virus (MDV) serotype I, serotype II, serotype III (type III is turkey herpesvirus), there can be illustrated the TK region (Ross L. et al., 1991), US10 region (Sakaguchi M. et al., 1994), the US2 region (Sondermeijer, P.j. et al., 1993), the region in between UL44 and UL45 and the region in between UL45 and UL46 (~~SEQ ID NO: 3~~ indicates a sequence from UL44 to UL46 (UVT-UL44-46 insertion site)) described in International Patent Publication W099/18215.

Please replace the paragraph beginning on page 14, line 32, with the following rewritten paragraph:

As the gene region nonessential for the growth of poxvirus, there can be used the TK gene region of fowlpox virus, pigeon poxvirus, quail pox (?), turkey poxvirus etc., the region in between the open reading frames (ORF) as described in U.S. Pat. No. 5,180,675 and Japanese Patent Publication No. 2766984, and the region described in U.S. Pat. No. 5,387,519. More specifically, an EcoRI fragment (7.3 kbp), an EcoRI-HindIII fragment (about 5.0 kbp), a BamHI fragment (about 4.0 kbp), and a HindIII fragment (about 5.2 kbp) derived from pigeon pox described in U.S. Pat. No. 5,387,519, and SpeI-HpaI (3026 bp) (~~SEQ ID NO: 4~~) (FPV-29 insertion site) in an EcoRI fragment (7.3 kbp).

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

Please replace the paragraph beginning on page 20, line 14, with the following rewritten paragraph:

There are four N-glycosylation sites in the TTM-1 portion of the amino acid sequence as set forth in SEQ ID NO: 12 of a plasmid pNZ40K-S described in International Patent Publication W097/36924 in which a MDVgB signal-encoding DNA has been ligated to the N-terminal of the antigen gene TTM-1 derived from Mycoplasma gallisepticum. Since there are no N-glycosylation sites in the region from the EcoRI site, the start of the TTM-1 portion, to the BglII site 83 bp downstream, the portion downstream of BglII was modified based on the above principle of alteration A to obtain a plasmid pGTPs40KS-Ngly having an modified TTM-1 gene in which the sequence downstream to the BglII site has the nucleotide sequence ~~as set forth in SEQ ID NO: 24~~ (TTM-1 portion of the modified pNZ40K-S (downstream of BglII)) that corresponds to the amino acid sequence ~~as set forth in SEQ ID NO: 23~~. The specific procedure to obtain this plasmid is as follows:

Please replace the six paragraphs beginning on page 21, line 2, with the following rewritten paragraphs:

First, PCR was performed with the primer 40KG-1 (SEQ ID NO: 13) and the primer 40KG-2R (SEQ ID NO: 14) to obtain a 136 bp fragment.

Similarly, PCR was performed with the primer 40KG-2 (SEQ ID NO: 15) and the primer 40 KG-3R (SEQ ID NO: 16) to obtain a 341 bp fragment.

Similarly, PCR was performed with the primer 40KG-3A (SEQ ID NO: 17) and the primer 40KG-4RA (SEQ ID NO: 18) to obtain 190 bp fragment.

Similarly, PCR was performed with the primer 40KG-4 (SEQ ID NO: 19) and the primer 40KG-5R

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

(~~SEQ ID NO: 20~~) to obtain a 359 bp fragment.

Similarly, PCR was performed with the primer 40KG-5 (~~SEQ ID NO: 21~~) and the primer 40KF-6R (~~SEQ ID NO: 22~~) to obtain a 218 bp fragment.

In the next PCR, using three fragments of a 136 bp RCE product of the primers 40KG-1 and 40KG-2R, a 341 bp PCR product of the primers 40KG-2 and 40KG-3R, and a 190 bp PCR product of the primers 40KG-3A and 40KG-4RA as the templates, PCR was performed in the above-mentioned condition with the primer 40KG-1 (~~SEQ ID NO: 13~~) and the primer 4KG-4RA (~~SEQ ID NO: 18~~) to obtain a 595 bp fragment.

Please replace the paragraph beginning on page 21, line 25, with the following rewritten paragraph:

Similarly, using a 359 bp PCR product of the primers 40KG-4 and 40KG-5R and a 218 bp PCR product of the primers 40KG-5 and 40KG-6R as the templates, PCR was performed in the above-mentioned condition with the primer 40KG-4 (~~SEQ ID NO: 19~~) and the primer 4KG-6R (~~SEQ ID NO: 22~~) to obtain a 539 bp fragment.

Please replace the paragraph beginning on page 21, line 31, with the following rewritten paragraph:

Furthermore, using a 595 bp PCR product of the primers 40KG-1 and 40KG-4RA and a 539 bp PCR product of the primers 40KG-4 and 40KG-6R as the templates, PCR was performed in the above-mentioned condition with the primer 40KG-1 (~~SEQ ID NO: 13~~) and the primer 4KG-6R (~~SEQ ID NO: 22~~) to obtain a partial fragment (1088 bp) of the modified TTM-1.

Please replace the paragraph beginning on page 22, line 8, with the following rewritten paragraph:

The amino acid sequence as set forth in SEQ ID NO: 25 4 encoded by the *mgc3* gene (GeneBank accession No. AB023292) as set forth in SEQ ID NO: 2 has 16 N-glycosylation sites. Among the 16 sites, since a site at the most 5' upstream side is replaced with the signal sequence at a later treatment, it was excluded from the target of alteration. For the remaining 15 sites, the base sequence was modified based on the principle of alteration A to obtain the plasmid pM11BTR containing the *mgc3* gene that has the nucleotide sequence as set forth in SEQ ID NO: 79 (modified *mgc3* gene (M11-BTR)) corresponding to the amino acid sequence as set forth in SEQ ID NO: 78 (modified MG3g antigen (M11-BTR)).

Please replace the six paragraphs on page 23, line 3, with the following rewritten paragraphs:

Using pUC-MGC3 as the template, PCR was performed with the primer M11-B (SEQ ID NO: 26) and the primer M11-2R (SEQ ID NO: 27) to obtain a 136 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-2 (SEQ ID NO: 28) and the primer M11-3R (SEQ ID NO: 29) to obtain a 92 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer Mll-3 (SEQ ID NO: 30) and the primer Mll-4RB (SEQ ID NO: 31) to obtain a 271 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-4B (SEQ ID NO: 32) and the primer M11-5R (SEQ ID NO: 33) to obtain a 116 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-5A (SEQ ID NO: 34) and the primer M11-7RA (SEQ ID NO: 35) to obtain a 439 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-7 (~~SEQ ID NO: 36~~) and the primer M11-2KR (~~SEQ ID NO: 37~~) to obtain a 201 bp fragment.

Please replace the paragraphs beginning on page 23, line 21, through page 26, line 33, as follows:

In the next PCR, using a 136 bp PCR product of the primer M11-B and the primer M11-2R and a 92 bp PCR product of the primer M11-2 and the primer M11-3R as the templates, PCR was performed with the primer M11-B (~~SEQ ID NO: 26~~) and the primer M11-3R (~~SEQ ID NO: 29~~) to obtain a 199 bp fragment.

Similarly, using a 271 bp PCR product of the primer M11-3 and the primer M11-4RB and a 116 bp PCR product of the primer M11-4B and the primer M11-5R as the templates, PCR was performed with the primer M11-3 (~~SEQ ID NO: 30~~) and the primer M11-5R (~~SEQ ID NO: 33~~) to obtain a 344 bp fragment.

Similarly, using a 439 bp PCR product of the primer M11-5A and the primer M11-7RA and a 201 bp PCR product of the primer M11-7 and the primer M11-KR as the templates, PCR was performed with the primer M11-5A (~~SEQ ID NO: 34~~) and the primer M11-KR (~~SEQ ID NO: 37~~) to obtain a 610 bp fragment.

Using these three PCR products, a PCR product (199 bp) of the primer M11-B and the primer M11-3R, and a PCR product (361 bp) of the primer M11-3 and the primer M11-5R, and a PCR product (610 bp) of the primer M11-5A and M11-KR as the templates, PCR was performed with the primer M11-B (~~SEQ ID NO: 26~~) and the primer M11-KR (~~SEQ ID NO: 37~~) to obtain a 1094 bp fragment. A 1070 bp fragment obtained by cleaving this PCR product (1094 bp) 10 of the primers M11-B and M11-KR with EcoRI and KpnI was inserted into a 2678 bp fragment of the plasmid pUC18 cleaved with EcoRI and KpnI to construct pM11BKR.

After analyzing the nucleotide sequence of this pM11BKR, a sequence different from the *mgc3* gene (GeneBank accession No. AB023292) registered in GeneBank was found. Furthermore, the sequence of the

original plasmid pUC-MGC3 was compared to confirm that it was not an error in PCR. As a result, it was confirmed that G at position 308 of SEQ ID NO: 2, a sequence registered at GeneBank, is T, G at position 311 is C, C at position 561 is G, and G at position 749 is T. It was demonstrated that due to an error in the base at position 561, the amino acid sequence encoded in this region is not N-Asn(N)-Gln(Q)-Thr(T) corresponding to the N-glycosylation site but Gln(Q)-Gln(Q)-Thr(T).

(2) Mutation of the KXR region (construction of pM11KXR)

Using pUC-MGC3 as the template, PCR was performed with the primer M11-K (~~SEQ ID NO: 38~~) and the primer M11-8R (~~SEQ ID NO: 39~~) to obtain a 151 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-8 (~~SEQ ID NO: 40~~) and the primer M11-10R (~~SEQ ID NO: 41~~) to obtain a 109 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-10 (~~SEQ ID NO: 42~~) and the primer M11-12RA (~~SEQ ID NO: 43~~) to obtain a 416 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-12A (~~SEQ ID NO: 44~~) and the primer M11-XR (~~SEQ ID NO: 45~~) to obtain a 350 bp fragment.

In the next PCR, using a 109 bp PCR product of the primer M11-8 and the primer M11-10R and a 416 bp PCR product of the primer M11-10 and the primer M11-12RA as the templates, PCR was performed with the primer M11-8 (~~SEQ ID NO: 40~~) and the primer M11-12RA (~~SEQ ID NO: 43~~) to obtain a 487 bp fragment.

Furthermore, using a 151 bp PCR product of the primer M11-K and the primer M11-8R and a 487 bp PCR product of the primer M11-8 and the primer M11-12RA as the templates, PCR was performed with the primer M11-K (~~SEQ ID NO: 38~~) and the primer M11-12RA (~~SEQ ID NO: 43~~) to obtain a 596 bp fragment.

Using the above two PCR products, a 596 bp PCR product of the primer M11-K and the primer M11-

12RA and a 350 bp PCR product of the primer M11-12A and the primer M11-XR as the templates, PCR was performed with the primer M11-K (SEQ ID NO: 38) and the primer M11-XR (SEQ ID NO: 45) to obtain a 908 bp fragment.

A 885 bp fragment obtained by cleaving this PCR product (908 bp) of the primer M11-K and the primer M11-XR with KpnI and XbaI was inserted into the plasmid pUC18 cleaved with KpnI and XbaI to construct pM11KXR.

After analyzing the nucleotide sequence of this pM11KXR, a sequence different from the *mgc3* gene (GeneBank accession No. AB023292) registered in GeneBank was found. Furthermore, the sequence of the original plasmid pUC-MGC3 was compared to confirm that it was not an error in PCR. As a result, it was confirmed that G at position 1279 of SEQ ID NO: 2, a sequence registered at GeneBank, is A, T at position 1729 is G, and C at position 1732 is G.

(3) Mutation of the XGTR region (construction of pM11XGTR)

Using pUC-MGC3 as the template, PCR was performed with the primer M11-XA (SEQ ID NO: 46) and the primer M11-13RA (SEQ ID NO: 47) to obtain a 238 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-13A (SEQ ID NO: 48) and the primer M11-14RA (SEQ ID NO: 49) to obtain a 266 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-14A (SEQ ID NO: 50) and the primer M11-15RA (SEQ ID NO: 51) to obtain a 168 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-15A (SEQ ID NO: 52) and the primer M11-16RA (SEQ ID NO: 53) to obtain a 123 bp fragment.

Using pUC-MGC3 as the template, PCR was performed with the primer M11-16A (SEQ ID NO: 54) and the primer M11-GTR (SEQ ID NO: 55) to obtain a 556 bp fragment.

In the next PCR, using a 238 bp PCR product of the primer M11-XA and the primer M11-13RA and a 266 bp PCR product of the primer M11-13A and the primer M11-14RA as the templates, PCR was performed with the primer M11-XA (~~SEQ ID NO: 46~~) and the primer M11-14RA (~~SEQ ID NO: 49~~) to obtain a 463 bp fragment.

Furthermore, using a 168 bp PCR product of the primer M11-14A and the primer M11-15RA and a 123 bp PCR product of the primer M11-15A and the primer M11-16RA as the templates, PCR was performed with the primer M11-14A (~~SEQ ID NO: 50~~) and the primer M11-16RA (~~SEQ ID NO: 53~~) to obtain a 253 bp fragment.

Using the above three fragments of a 463 bp PCR product of the primer M11-XA and the primer M11-14RA, a 253 bp PCR product of the primer M11-14A and the primer M11-16RA, and a 556 bp PCR product of the primer M11-16A and the primer M11-GTR as the templates, PCR was performed with the primer M11-XA (~~SEQ ID NO: 46~~) and the primer M11-GTR (~~SEQ ID NO: 55~~) to obtain a 1192 bp fragment.

Please replace the paragraphs beginning on page 27, line 22, as follows:

A 2676 bp fragment obtained by cleaving, with HindIII and SalI pUC18XG described in International Patent Publication W099/18215 in which the restriction sites of pUC18 were modified and an annealing product of a 5'-end-phosphorylated linker H'-S-H-S-P-S1 (~~SEQ ID NO: 56~~) and a linker H'-S-H-S-P-S2 (~~SEQ ID NO: 57~~) were ligated to construct pUCSfi-H-S.

(2) Construction of pGHPs

Three fragments, a 2661 bp fragment obtained by cleaving pUCSfi-H-S with HindIII and EcoRI, an annealing product of a 5'-end-phosphorylated linker S-B-E1 (~~SEQ ID NO: 58~~) and a linker S-B-E2 (~~SEQ ID NO: 59~~), and a 137 bp fragment obtained by cleaving, with HindIII and SalI, the plasmid described in

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

International Patent Publication WO97/36924 containing the poxvirus late and early promoter, were ligated to construct a plasmid pGHPs.

Please replace the paragraphs beginning on page 29, line 32, through page 30, line 15, as follows:

Using pNZ29RMG40KM11CS-G constructed in Example 4(2) as the template, PCR was performed with the primer M11-Sfi (~~SEQ ID NO: 60~~) and the primer M11-5RB (~~SEQ ID NO: 61~~) in the condition described in Example 1 to obtain a 836 bp fragment.

Using pNZ29RMG40KM11CS-G constructed in Example 4(2) as the template, PCR was performed with the primer M11-5C (~~SEQ ID NO: 62~~) and the primer M11-KRA (~~SEQ ID NO: 63~~) in a similar condition to obtain a 618 bp fragment.

Using these two PCR products as the templates, PCR was performed with the primer M11-Sfi (~~SEQ ID NO: 60~~) and the primer M11-KRA (~~SEQ ID NO: 63~~) to obtain a 1400 bp fragment. A 1368 bp fragment obtained by cleaving this 1400 bp PCR product with SfiI and KpnI and a 9032 bp fragment obtained by cleaving pNZ29RMG40KM11CS-G constructed in Example 4(2) with SfiI and KpnI were ligated to construct a plasmid pNZ29RMG40KM11HCS-G2 having the modified MDVgB signal DNA, the modified TTM-1 gene, and the re-modified mgc3 gene.

Please replace the paragraphs beginning on page 30, line 34, through page 32, line 1, as follows:

Using pGIPEC described in Japanese Unexamined Patent Publication (Kokai) No. 2001-188 as the template, PCR was performed in the condition created in Example 1 with the primer pCMV-1 (~~SEQ ID NO: 64~~) and the primer pPecIR (~~SEQ ID NO: 65~~) to obtain a 293 bp fragment.

Preliminary Amendment
Takeshi OKUDA et al.

U.S. Patent Application S.N. 09/901,572
Attorney Docket No. 010898

of MDV-1 was deleted in pGHCMV(-).

Please replace the paragraphs beginning on page 32, line 24, through page 33, line 11, as follows:

Using pBK-CMV (Stratagene) as the template, PCR was performed with the primer PolyA-SfiF2 (~~SEQ ID NO: 74~~) and the primer PolyA-SalKpn (~~SEQ ID NO: 75~~) to obtain a 313 bp fragment having the PolyA signal of SV40. A 297 bp fragment obtained by cleaving this 313 bp fragment with SfiI and KpnI and a 4222 bp fragment obtained by cleaving pBac(dHS) obtained in the above (9) with SfiI and KpnI were ligated to construct pGIBacpA2nd.

(11) Construction of pGIBac40KS2nd

Using pGTPs40K-S described in International Patent Publication W097/36924 as the template, PCR was performed with the primer 40KS-B (~~SEQ ID NO: 76~~) and 40KG-6R (~~SEQ ID NO: 22~~) to obtain a 1359 bp fragment. A 1345 bp fragment obtained by cleaving this 1359 bp fragment with BamHI and SalI and a 4222 bp fragment obtained by cleaving pGIBacpA2nd obtained in the above (10) with BamHI and SalI were ligated to construct a plasmid pGIBac40KS2nd.

(12) Construction of p45/46Bac40KS+2nd

A 3169 bp fragment obtained by cleaving pGIBac40KS2nd obtained in the above (11) with BglI was inserted into the SfiI site of pNZ45/46Sfi having the UL44 to UL46 region (~~SEQ ID NO: 3~~) (HVT-UL44-46 insertion site) of HVT described in W097/36924 to construct a plasmid p45/46Bac40KS+2nd having the TTM-1 gene.

Please replace the paragraph beginning on page 33, line 33, with the following rewritten

paragraph:

Using pUTTM-1 described in U.S. Pat. No. 5,489,430 as the template, PCR was performed with primer pMG40K-1 (~~SEQ ID NO: 77~~) and the primer 40KG-6R (~~SEQ ID NO: 22~~) to obtain a 1259 fragment. A 1237 bp fragment obtained by cleaving this 1259 bp fragment with BamHI and SalI and a 7317 bp fragment obtained by cleaving pNZ45/46BacpA2nd obtained in the above (15) with BamHI and SalI were ligated to construct a plasmid pNZ45/46Bac40KpA+2nd.

Please amend pages 55, line 1 through page 59, line 4, as follows:

Sequence listing Free text

- | | |
|------------------------------|-----------------------------------------------------------------|
| Sequence No. 1 | TTM-1 gene (behind EcoRI) |
| Sequence No. 2 | mgc3 gene |
| Sequence No. 3 | HVT-UL44-46 insertion site |
| Sequence No. 4 | FPV-29 insertion site |
| Sequence No. 5 | MDVgB signal (amino acid) |
| Sequence No. 6 | modified MDVgB signal (amino acid) |
| Sequence No. 7 | modified MDVgB signal (gene) |
| Sequence No. 8 | Rabies virus glycoprotein G (gG) signal (amino acid) |
| Sequence No. 9 | synthetic DNA-1: 1BN for Rabies gG signal |
| Sequence No. 10 | synthetic DNA-2: 1BC for Rabies gG signal |
| Sequence No. 11 | modified Rabies virus glycoprotein G signal |
| Sequence No. 12 3 | TTM-1 portion of pNZ40K-S (amino acid) |
| Sequence No. 13 | synthetic DNA (primer): 40KG-1 (23 mer) |

- Sequence No. 14 — synthetic DNA (primer): 40KG-2R (22 mer)
- Sequence No. 15 — synthetic DNA (primer): 40KG-2 (23 mer)
- Sequence No. 16 — synthetic DNA (primer): 40KG-3R (21 mer)
- Sequence No. 17 — synthetic DNA (primer): 40KG-3A (36 mer)
- Sequence No. 18 — synthetic DNA (primer): 40KG-4RA (34 mer)
- Sequence No. 19 — synthetic DNA (primer): 40KG-4 (24 mer)
- Sequence No. 20 — synthetic DNA (primer): 40KG5R (28 mer)
- Sequence No. 21 — synthetic DNA (primer): 40KG-5 (27 mer)
- Sequence No. 22 — synthetic DNA (primer): 40KG-6R (37 mer)
- Sequence No. 23 — TTM-1 portion of modified pNZ40K-S (downstream of BflI) (amino acid)
- Sequence No. 24 — TTM-1 portion of the modified pNZ40K-S (downstream of BflI) (gene)
- Sequence No. 25 4 — MGC3 protein encoded by the mgc3 gene (amino acid)
- Sequence No. 26 — synthetic DNA (primer): M11-B (26 mer)
- Sequence No. 27 — synthetic DNA (primer): M11-2R (20 mer)
- Sequence No. 28 — synthetic DNA (primer): M11-2 (20 mer)
- Sequence No. 29 — synthetic DNA (primer): M11-3R (20 mer)
- Sequence No. 30 — synthetic DNA (primer): M11-3 (20 mer))
- Sequence No. 31 — synthetic DNA (primer): M11-4RB (30 mer)
- Sequence No. 32 — synthetic DNA (primer): M11-4B (30 mer)
- Sequence No. 33 — synthetic DNA (primer): M11-5R (20 mer)
- Sequence No. 34 — synthetic DNA (primer): M11-5A (30mer)
- Sequence No. 35 — synthetic DNA (primer): M11-7RA (20 mer)

Sequence No. 36 — synthetic DNA (primer): M11-7 (20 mer)
Sequence No. 37 — synthetic DNA (primer): M11-KR (20 mer)
Sequence No. 38 — synthetic DNA (primer): M11-K (20 mer)
Sequence No. 39 — synthetic DNA (primer): M11-8R (30 mer)
Sequence No. 40 — synthetic DNA (primer): M11-8 (30 mer)
Sequence No. 41 — synthetic DNA (primer): M11-10R (25 mer)
Sequence No. 42 — synthetic DNA (primer): M11-10 (25 mer)
Sequence No. 43 — synthetic DNA (primer): M11-12RA (30 mer)
Sequence No. 44 — synthetic DNA (primer): M11-12A (30 mer)
Sequence No. 45 — synthetic DNA (primer): M11-XR (20 mer)
Sequence No. 46 — synthetic DNA (primer): M11-XA (30 mer)
Sequence No. 47 — synthetic DNA (primer): M11-13RA (30 mer)
Sequence No. 48 — synthetic DNA (primer): M11-13A (30 mer)
Sequence No. 49 — synthetic DNA (primer): M11-14RA (30 mer)
Sequence No. 50 — synthetic DNA (primer): M11-14A (30 mer)
Sequence No. 51 — synthetic DNA (primer): M11-15RA (30 mer)
Sequence No. 52 — synthetic DNA (primer): M11-15A (30 mer)
Sequence No. 53 — synthetic DNA (primer): M11-16RA (30 mer)
Sequence No. 54 — synthetic DNA (primer): M11-16A (30 mer)
Sequence No. 55 — synthetic DNA (primer): M11-GTR (30 mer)
Sequence No. 56 — synthetic DNA (linker): IP-S-H-S-P-S1 (36 mer)
Sequence No. 57 — synthetic DNA (linker): IP-S-H-S-S2 (36 mer)

- Sequence No. 58 ——— synthetic DNA (linker): S-B-E1 (35-mer)
- Sequence No. 59 ——— synthetic DNA (linker): S-B-E2 (35-mer)
- Sequence No. 60 ——— synthetic DNA (primer): M11-Sfi (30-mer)
- Sequence No. 61 ——— synthetic DNA (primer): M11-5RB (30-mer)
- Sequence No. 62 ——— synthetic DNA (primer): M11-5C (30-mer)
- Sequence No. 53 ——— synthetic DNA (primer): M11-KRA (30-mer)
- Sequence No. 64 ——— synthetic DNA (primer): pCMV-1 (30-mer)
- Sequence No. 65 ——— synthetic DNA (primer): pPec1R (30-mer)
- Sequence No. 66 ——— synthetic DNA (primer): pCMV-01 (30-mer)
- Sequence No. 67 ——— synthetic DNA (primer): pCMV-R1 (30-mer)
- Sequence No. 68 ——— synthetic DNA (primer): pGHP-F (29-mer)
- Sequence No. 69 ——— synthetic DNA (primer): pGHP-R (30-mer)
- Sequence No. 70 ——— synthetic DNA (linker): Linker 1 (23-mer)
- Sequence No. 71 ——— synthetic DNA (linker): Linker 2 (23-mer)
- Sequence No. 72 ——— synthetic DNA (linker): Linker 3 (16-mer)
- Sequence No. 73 ——— synthetic DNA (linker): Linker 4 (12-mer)
- Sequence No. 74 ——— synthetic DNA (primer): PolyA-SfiF2 (30-mer)
- Sequence No. 75 ——— synthetic DNA (primer): PolyA-San1Kpn (30-mer)
- Sequence No. 76 ——— synthetic DNA (primer): 40KS-B (22-mer)
- Sequence No. 77 ——— synthetic DNA (primer): pMG40K-1 (30-mer)
- Sequence No. 78 ——— modified MGC3-antigen (M11-BTR) (amino acid)
- Sequence No. 79 ——— modified mgc3-gene (M11-BTR) (DNA)